

THE DOUBLY HEAVY BARYONS IN THE NONPERTURBATIVE QCD APPROACH

I.M.NARODETSKII AND M.A.TRUSOV

ITEP, Moscow, Russia
E-mail: naro@heron.itep.ru

We present some piloting calculations of the masses of the doubly heavy baryons in the framework of the simple approximation within the nonperturbative string approach. The simple analytical results for dynamical masses of heavy and light quarks and eigenvalues of the effective QCD Hamiltonian are presented.

The purpose of this talk is to present the results of the calculation ¹ of the masses and wave functions of the heavy baryons in a simple approximation within the nonperturbative QCD (see ² and references therein). The starting point of the approach is the Feynman-Schwinger representation for the three quark Green function in QCD in which the role of the time parameter along the trajectory of each quark is played by the Fock-Schwinger proper time. The proper and real times for each quark related via a new quantity that eventually plays the role of the dynamical quark mass. The final result is the derivation of the Effective Hamiltonian, see Eq. (1) below. In contrast to the standard approach of the constituent quark model the dynamical mass m_i is not a free parameter but it is expressed in terms of the current mass $m_i^{(0)}$ defined at the appropriate scale of $\mu \sim 1$ GeV from the condition of the minimum of the baryon mass M_B as function of m_i : $\frac{\partial M_B(m_i)}{\partial m_i} = 0$. Technically, this has been done using the einbein (auxiliary fields) approach, which is proven to be rather accurate in various calculations for relativistic systems.

This method was already applied to study baryon Regge trajectories ³ and very recently for computation of magnetic moments of light baryons ⁴. The essential point of this talk is that it is very reasonable that the same method should also hold for hadrons containing heavy quarks. As in ⁴ we take as the universal parameter the QCD string tension σ fixed in experiment by the meson and baryon Regge slopes. We also include the perturbative Coulomb interaction with the frozen coupling $\alpha_s(1 \text{ GeV}) = 0.4$.

Consider the ground state baryons without radial and orbital excitations in which case tensor and spin-orbit forces do not contribute perturbatively.

5 survives in the perturbative approximation. The EH has the following form

$$H = \sum_{i=1}^3 \left(\frac{m_i^{(0)2}}{2m_i} + \frac{m_i}{2} \right) + H_0 + V, \quad (1)$$

where H_0 is the non-relativistic kinetic energy operator, $m_i^{(0)}$ are the current quark masses and m_i are the dynamical quark masses to be found from the minimum condition, and V is the sum of the perturbative one gluon exchange potential V_c and the string potential V_{string} . The string potential has been calculated in ³ as the static energy of the three heavy quarks: $V_{string}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = \sigma R_{min}$, where R_{min} is the sum of the three distances $|\mathbf{r}_i|$ from the string junction point, which for simplicity is chosen as coinciding with the center-of-mass coordinate.

We use the hyper radial approximation (HRA) in the hyper-spherical formalism approach. In the HRA the three quark wave function depends only on the hyper-radius $R^2 = \boldsymbol{\rho}^2 + \boldsymbol{\lambda}^2$, where $\boldsymbol{\rho}$ and $\boldsymbol{\lambda}$ are the three-body Jacobi variables: $\boldsymbol{\rho}_{ij} = \sqrt{\frac{\mu_{ij}}{\mu}}(\mathbf{r}_i - \mathbf{r}_j)$, $\boldsymbol{\lambda}_{ij} = \sqrt{\frac{\mu_{ij,k}}{\mu}}\left(\frac{m_i\mathbf{r}_i + m_j\mathbf{r}_j}{m_i + m_j} - \mathbf{r}_k\right)$, where $\mu_{ij} = \frac{m_i m_j}{m_i + m_j}$, $\mu_{ij,k} = \frac{(m_i + m_j)m_k}{m_i + m_j + m_k}$, and μ is an arbitrary parameter with the dimension of mass which drops off in the final expressions. Introducing the reduced function $\chi(R) = R^{5/2}\psi(R)$ and averaging $V = V_c + V_{string}$ over the six-dimensional sphere one obtains the Schrödinger equation

$$\frac{d^2\chi(R)}{dR^2} + 2\mu \left[E_n + \frac{a}{R} - bR - \frac{15}{8\mu R^2} \right] \chi(R) = 0, \quad (2)$$

where

$$a = \frac{2\alpha_s}{3} \cdot \frac{16}{3\pi} \sum_{i<j} \sqrt{\frac{\mu_{ij}}{\mu}}, \quad b = \sigma \cdot \frac{32}{15\pi} \sum_{i<j} \sqrt{\frac{\mu(m_i + m_j)}{m_k(m_1 + m_2 + m_3)}}, \quad (3)$$

We use the same parameters as in Ref. ⁵: $\sigma = 0.17$ GeV, $\alpha_s = 0.4$, $m_q^{(0)} = 0.009$ GeV, $m_s^{(0)} = 0.17$ GeV, $m_c^{(0)} = 1.4$ GeV, and $m_b^{(0)} = 4.8$ GeV. We solve Eq. (2) by the variational method introducing a simple variational Ansatz $\chi(R) \sim R^{5/2}e^{-\mu p^2 R^2}$, where p is the variational parameter. Then the three-quark Hamiltonian admits explicit solutions for the energy and the ground state eigenfunction: $E \approx \min_p E(p)$, where

$$E(p) = \langle \chi | H | \chi \rangle = 3p^2 - (a\sqrt{\mu}) \cdot \frac{3}{4} \sqrt{\frac{\pi}{2}} \cdot p + (b/\sqrt{\mu}) \cdot \frac{15}{16} \sqrt{\frac{\pi}{2}} \cdot p^{-1}. \quad (4)$$

Table 1. Comparison of results of analytical and numerical variational calculations for Λ_b and Λ_c baryons (all quantities are in units of GeV)

Baryon	Λ_b		Λ_c	
	Numerical calculation	Analytical calculation	Numerical calculation	Analytical calculation
E_0	1.06	1.08	1.18	1.16
m_q	0.56	0.56	0.52	0.53
m_Q	4.84	4.82	1.50	1.47

The dynamical masses m_i and the ground state eigenvalues E_0 are given for various baryons in Table 1 of Ref. ¹. The dynamical values of light quark mass $m_q \sim \sqrt{\sigma} \sim 450 - 500$ MeV ($q = u, d, s$) qualitatively agree with the results of Ref. ⁵ obtained from the analysis of the heavy-light ground state mesons. For the heavy quarks ($Q = c$ and b) the variation in the values of their dynamical masses m_Q is marginal. This is illustrated by the simple analytical results for Qud baryons. These results were obtained from the approximate solution of equation $\left. \frac{dE}{dp} \right|_{p=p_0} = 0$ in the form of expansion in the small parameters $\xi = \sqrt{\sigma}/m_Q^{(0)}$ and α_s . Omitting the intermediate steps one has

$$\begin{aligned}
E_0 &= 3\sqrt{\sigma} \left(\frac{6}{\pi} \right)^{1/4} \left(1 + A \cdot \xi - \frac{5}{3} B \cdot \alpha_s + \dots \right) \\
m_q &= \sqrt{\sigma} \left(\frac{6}{\pi} \right)^{1/4} (1 - A \cdot \xi + B \cdot \alpha_s + \dots), \\
m_Q &= m_Q^{(0)} (1 + 2A \cdot \xi^2 + \dots)
\end{aligned}$$

where for our variational Ansatz $A = \frac{\sqrt{2}-1}{2} \left(\frac{6}{\pi} \right)^{1/4}$, $B = \frac{4+\sqrt{2}}{18} \sqrt{\frac{6}{\pi}}$. Accuracy of this approximation is illustrated in Table 1.

To calculate hadron masses we, as in Ref. ³, first renormalize the string potential: $V_{\text{string}} \rightarrow V_{\text{string}} + \sum_i C_i$, where the constants C_i take into account the residual self-energy (RSE) of quarks. In what follows we adjust the RSE constants C_i to reproduce the center-of-gravity for baryons with a given flavor. As a result we obtain $C_q = 0.34$, $C_s = 0.19$, $C_c \sim C_b \sim 0$.

We keep these parameters fixed to calculate the masses given in Table 2, namely the spin-averaged masses (computed without the spin-spin term) of the lowest double heavy baryons. In this Table we also compare our pre-

Table 2. Masses of baryons containing two heavy quarks

State	present work	Ref. ⁶	Ref. ⁷	Ref. ⁸	Ref. ⁹
$\Xi\{qcc\}$	3.69	3.70	3.71	3.66	3.48
$\Omega\{scc\}$	3.86	3.80	3.76	3.74	3.58
$\Xi\{qcb\}$	6.96	6.99	6.95	7.04	6.82
$\Omega\{scb\}$	7.13	7.07	7.05	7.09	6.92
$\Xi\{qbb\}$	10.16	10.24	10.23	10.24	10.09
$\Omega\{sbb\}$	10.34	10.30	10.32	10.37	10.19

dictions with the results obtained using the additive non-relativistic quark model with the power-law potential ⁶, relativistic quasipotential quark model ⁷, the Feynman-Hellmann theorem ⁸ and with the predictions obtained in the approximation of double heavy diquark ⁹.

In conclusion, we have employed the general formalism for the baryons, which is based on nonperturbative QCD and where the only inputs are σ , α_s and two additive constants, C_q and C_s , the residual self-energies of the light quarks. Using this formalism we have also performed the calculations of the spin-averaged masses of baryons with two heavy quarks. One can see from Table 2 that our predictions are especially close to those obtained in Ref. ⁶ using a variant of the power-law potential adjusted to fit ground state baryons.

Acknowledgements

This work was supported in part by RFBR grants ## 00-02-16363 and 00-15-96786.

References

1. I.M.Narodetskii and M.A.Trusov, Yad. Fiz., **65**, in press [hep-ph/0104019]
2. Yu.A.Simonov, Lectures given at the XVII International School of Physics "QCD: Perturbative or Nonperturbative", Lisbon 1999 [hep-ph/9911237]
3. M.Fabre de la Ripelle and Yu.A.Simonov, Ann. Phys. (N.Y.) **212**, 235 (1991).
4. B.O.Kerbikov, Yu.A.Simonov, Phys. Rev. **D62**, 093016 (2000).
5. Yu.S.Kalashnikova and A.Nefediev, Phys. Lett. B **492**, 91 (2000).
6. E.Bagan *et al.* Z.Phys. C **64**, 57 (1994).

7. D.Ebert *et al.*, Z. Phys. C **76**, 111 (1997).
8. R.Roncaglia *et al.*, Phys. Rev. D **52**, 1248 (1995).
9. A.K.Likhoded and A.I.Onishchenko, hep-ph/9912425